

Anti-Tau [pS422] Mouse Monoclonal Antibody (clone 2H9)

Ref. 4BDX-1501S

Biomolecule

Anti-Tau [pS422]
Mouse monoclonal antibody

Clone

2H9

Size

20 µg/20 µL

Formulation

Solution in PBS at 1 mg/mL

Storage

+4°C / -20°C

Immunogen

Peptide

Specificity

Anti-Tau phosphorylated
at serine 422

Cross-reactivity

Human, Mouse, Rat

Immunoglobulin type

Human Tau specific
mouse IgG

Isotype

IgG2a Kappa

Applications

Western-blot,
Immunohistochemistry,
Immunofluorescence,
Immunotherapy

• **Preparation**

This antibody was produced from a mouse hybridoma resulting from a mouse immunized with a peptide covering the human Tau protein sequence with the phospho-serine 422.

• **Purity**

Mouse monoclonal antibodies 2H9 was purified by protein A/G affinity chromatography. Purity > 90%, as determined by SDS-PAGE and visualized by Coomassie blue staining.

• **Concentration**

The measured concentration of the purified Anti-Tau [pS422] was 1 mg/ml as determined using a total protein concentration assay.

• **Specificity**

Determined by its ability to recognise human Tau phosphorylated at serine 422.

• **Storage**

Store at +4°C for short term use (1-2 weeks) - Store at -20°C for long term use.

• **Applications**

Recommended concentration of use are:

Western-blot: 0.2 µg/mL (working with cell and tissue).

IHC/IF: 2 µg/mL (working with cell, tissue, frozen and paraffin embedded tissue).

• General information

Tau proteins are encoded by a single gene *MAPT* (Gene ID: 4137). They belong to the family of microtubule-associated proteins. The phosphorylation of Tau protein at serine 422 is a pathological epitope and a specific marker of neurofibrillary degeneration, one of the neuropathological hallmark of Alzheimer's disease and also found in more than 20 Tauopathies. On cerebral tissue, frozen or paraffin embedded tissue, 2H9 Tau antibody detects neurons in neurofibrillary degeneration. The serine 422 is phosphorylated by the following kinases: CaMKII (Calcium Calmodulin-kinase II), PKA (protein kinase A), JNKs (c-Jun N-terminal kinases), ERK1/2 or p42/44 MAPK (mitogen-activated kinases), p38 MAPK and MARK (Microtubule-associated protein/microtubule affinity-regulating kinase).

• References


Troquier L, Caillierez R, Burnouf S, Fernandez-Gomez FJ, Grosjean ME, Zommer N, Sergeant N, Schraen-Maschke S, Blum D and Buee L (2012) Targeting phospho-Ser422 by active Tau Immunotherapy in the THY-Tau22 mouse model: a suitable therapeutic approach. *Curr Alzheimer Res.* 9(4):397-405.

Sergeant N, Bretteville A, Hamdane M, Caillet-Boudin ML, Grognet P, Bombois S, Blum D, Delacourte A, Pasquier F, Vanmechelen E, Schraen-Maschke S, and Buee L (2008) Biochemistry of Tau in Alzheimer's disease and related neurological disorders. *Expert review of proteomics* 5:207–224.


Sergeant N, Vingtdoux V, Eddarkaoui S, Gay M, Evrard C, Le Fur N, Laurent C, Caillierez R, Obriot H, Larchanché PE, Farce A, Coevoet M, Carato P, Kouach M, Descat A, Dallemagne P, Buée-Scherrer V, Blum D, Hamdane M, Buée L, Melnyk P. New piperazine multi-effect drugs prevent neurofibrillary degeneration and amyloid deposition, and preserve memory in animal models of Alzheimer's disease. *Neurobiol Dis.* 2019 Sep;129:217-233.

Tautou M, Eddarkaoui S, Descamps F, Larchanché PE, El Bakali J, Goveas LM, Dumoulin M, Lamarre C, Blum D, Buée L, Melnyk P, Sergeant N. A β -Secretase Modulator Decreases Tau Pathology and Preserves Short-Term Memory in a Mouse Model of Neurofibrillary Degeneration. *Front Pharmacol.* 2021 Jun 29;12:679335.


Lam S, Petit F, Herard AS, Boluda S, Eddarkaoui S, Guillermier M, The brain bank Neuro-CEB neuropathology network, Buee L, Duyckaerts C, Haik S, Picq JL, Dhenain M. Amyloid and tau pathologies, cognitives impairments and cerebral atrophy after Alzheimer brain extracts inoculation in a primate. *Acta Neuropath.* 2021 (in press).



Western-blotting of human brain tissue (1: control, 2 and 3: Alzheimer)



Immunohistochemistry of Thy-Tau22 transgenic model of neurofibrillary degeneration



Immunofluorescence of Thy-Tau22 transgenic model of neurofibrillary degeneration

TO PLACE AN ORDER



commercial@4biodx.com



www.4biodx.com



(33) 320 622 079



SPQI SAS
82 rue Jeanne d'Arc, 59000 Lille, France

RCS Lille Metropole 820 961 514 – Code APE 7219Z
VAT : FR 40 820 961 514 000 24