

Anti-ProAKAP4 Mouse Monoclonal Antibody (clone 6F12)

Ref. 4BDX-1701S

Biomolecule

Anti-proAKAP4 mouse monoclonal antibody

Clone 6F12

Size 20 μg in 20 μL

Formulation Solution in PBS at 1 mg/mL

Storage +4°C / -20°C

Immunogen Peptide

Specificity AKAP4 prodomain

Cross-reactivity Human, Horse, Bull, Mouse, Ram, Camel

Immunoglobulin type Human AKAP4 specific mouse IgG

Isotype IgG2a Kappa

Applications WB, IF, IHC

• Preparation

This antibody was produced from a mouse hybridoma resulting from a mouse immunized with a peptide covering the prodomain of human AKAP4 protein sequence (Uniprot ref. Q5JQC9) which is 70% homologous between mammals.

• <u>Purity</u>

Mouse monoclonal antibodies 6F12 were purified by protein A/G affinity chromatography. Purity > 90%, as determined by SDS-PAGE and visualized by silver staining.

<u>Concentration</u>

The measured concentration of the purified anti-proAKAP4 antibodies was 1mg/mL as determined using a total protein concentration assay.

• Specificity

Determined by its ability to recognize the prodomain of human AKAP4 protein. This monoclonal antibody (clone 6F12) only recognizes the proAKAP4 (110 kDa / 854 AA), and the prodomain (21 kDa) and does not react with the AKAP4 (82 kDa / 665 AA). The clone 6F12 reacts also with proAKAP4 proteins from horse, bull, mouse, rat and ram semen.

• Storage

Store at +4°C for short-term use (1-2 weeks) - Store at -20°C for long-term use.

• Applications

Recommended concentrations of use are: Western-blot: 0.1 μg/mL IHC / IF: 5 μg/mL

<u>General information</u>

Human AKAP4 (A-Kinase Anchor Protein 4) protein is encoded by a single gene located on chromosome X. The proAKAP4 polypeptide is converted into mature AKAP4 by proteolytic cleavage of the amino-terminal prodomain made of 188 amino acids. AKAP4



and its precursor proAKAP4 are both major components of the fibrous sheath of the sperm flagellum. AKAP4 protein belongs to the family of A-kinase anchor proteins (AKAPs) all sharing a common function of binding to the regulatory subunit of protein kinase A (PKA) and confining the PKA holoenzyme to discrete locations within the cell. AKAP4 is also named AKAP-4, AKAP82 (A-Kinase Anchor Protein 82 KDa), PRKA4 (Protein Kinase Anchoring Protein 4), HI, CT99 (Cancer/Testis Antigen 99), FSC1 (Fibrous sheath component 1) or P82. AKAP4 plays a major role in flagellum formation, structuration, sperm motility, capacitation, and fertility.

Main References

Dordas-Perpinyà M, Sergeant N, Ruelle I, Bruyas JF, Charreaux F, Michaud S, Carracedo S, Catalán J, Miró J, Delehedde M, Briand-Amirat L (2022) ProAKAP4 Semen Concentrations as a Valuable Marker Protein of Post-Thawed Semen Quality and Bull Fertility: A Retrospective Study. Vet Sci. May 6;9(5):224.

Malo C, Carracedo S, Delehedde M, Sergeant N and Skidmore L (2021) Identification of PROAKAP4 in Dromedary sperm and their correlation with monthly semen parameters. Reproduction & Fertility. Vol.2(4):268-279.

Riesco M, Anel-Lopez L, Neila-Montero M, Palacin-Martinez C, Montes-Garrido R, Alvarez M, de Paz P, Anel L (2020) ProAKAP4 as Novel Molecular Marker of Sperm Quality in Ram: An Integrative Study in Fresh, Cooled and Cryopreserved Sperm. Biomolecules. 10(7):1046. doi: 10.3390/biom10071046.

Delehedde M, Carracedo S, Selleslagh M, Eddarkaoui S, Amirat-Briand L and Sergeant N (2019) ProAKAP4 polypeptide as a biomarker of sperm functionality and male fertility disorders. Int J Gynecol and Reprod Sci. Vol. 2(1):13-19.

Blommaert D, Sergeant N, Delehedde M, Jouy N, Mitchell V, Franck T, Donnay I, Lejeune JP and Serteyn D (2019) Expression, localization, and concentration of A-kinase anchor protein 4 (AKAP4) and its precursor (proAKAP4) in equine semen: promising marker correlated to the total and progressive motility in thawed spermatozoa. Theriogenology. Vol. 131:52-60.


Sergeant N, Briand-Amirat L, Bencharif D and Delehedde M (2019) The sperm specific protein proAKAP4 as an innovative marker to evaluate sperm quality and fertility. Journal of Dairy & Veterinary Sciences. Vol. 11:01-19.

Application examples

The monoclonal antibody (clone 6F12) only recognizes the full-length of AKAP4 called proAKAP4 (110 kDa / 854 AA), the prodomain (21 kDa) and does not react with the AKAP4 (82 kDa / 665 AA).



Western-blotting of human sperm protein extracts



Immunofluorescence of human spermatozoa



RCS Lille Metropole 820 961 514 – Code APE 7219Z VAT : FR 40 820 961 514 000 24