



Anti-MCM7 antibody, rabbit polyclonal IgG

Key words: DNA replication licensing factor, MCM complex, DNA replication initiation, G1/S transition, DNA damage response, DNA helicase,

Function: MCM7 (human; 718 aa, 80 kDa) acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for S-phase checkpoint activation upon UV-induced damage.

Applications

- 1) Western blotting (1/1,000~1/5,000 dilution).
- 2) Immunoprecipitation (assay dependent)
- 3) Immunofluorescence staining (1/200~1/1,000 dilution).

Other applications have not been tested.

Immunogen: Purified His6-tagged human MCM7 protein encompassing 562 -719 amino acids.

Reactivity: Reacts with human mouse, rat and hamster. Not tested in other species.

Product: Purified IgG from the rabbit antiserum. 1 mg/ml in PBS, 50% glycerol, filter-sterilized. Azide- and carrier-free.

Size: 100 ug

Storage: Shipped at 4°C. Upon arrival, spin-down, aliquot and store at -20°C.

Data Link UniProtKB/Swiss-Prot [P33993](#) MCM7_HUMAN

References: This antibody was described and used in the following publications.

1. Fujita M et al. (1996) hCDC47, a human member of the MCM family. Dissociation of the nucleus-bound form during S phase. J Biol Chem. 271:4349-54. [PMID 8626784](#) [Free Article](#). **WB, IP, IF**
2. Fujita M. et al. (1997) In vivo interaction of human MCM heterohexameric complexes with chromatin. Possible involvement of ATP. J Biol Chem. 272:10928-35. [PMID 9099751](#) [Free Article](#). **WB, IP**
3. Fujita M. et al. (2002) Nuclear organization of DNA replication initiation proteins in mammalian cells. J Biol Chem. 277:10354-61. [PMID 11779870](#) [Free Article](#). **WB, IP, IF**.



Anti-MCM7 antibody




Fig. 1. Identification of MCM7 protein in whole cell extracts of human cells by western blotting using anti-MCM7 antibody.

Lane 1; SiHa cells

Lane 2; C33A cells

Lane 3; WI38 cells

All cell lines are cervical cancer derived. Samples are obtained from approximately 10^5 cells.




Fig. 2 Identification of MCM7 protein in whole cell extracts of human and mouse cells by western blotting using anti-MCM7 antibody..

Lane 1. Size marker proteins in kDa.

Lane 2. Extract of HeLa cells untreated (-).

Lane 3. Extract of HeLa cells treated with 100 nM adriamycin for 24 hr (+)

Lane 4. Extract of NIH3T3 (mouse) cells.

Anti-MCM7 antibody was used at 1/2,000 dilution.

* Indicates the band of MCM7 protein

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia, Inc.



COSMO BIO Co., LTD.
Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

http://www.cosmobio.co.jp/index_e.asp

E-mail: export@cosmobio.co.jp

Phone : +81-3-5632-9617

FAX : +81-3-5632-9618



Anti-MCM7 antibody




Fig. 3. Immunoprecipitation of MCM7 protein from crude extract of human fibroblast cell line WI38 by using anti-MCM7 antibody..

Lane 1; Immunoprecipitation with pre-immune serum

Lane 2; Immunoprecipitation with anti-MCM7 antiserum.

Cells were labeled with S^{35} methionine and MCM7 was immunoprecipitated with the anti-MCM7 antibody followed by SDS-PAGE and autoradiography.




Fig. 4. Immunofluorescence staining and confocal microscopic analysis of MCM7 in G_1 phase HeLa cell nucleus by using anti-MCM7 antibody after treatment with protein cross-linking reagent, DSP and chromatin extraction. The processed cells were fixed with formaldehyde before staining.

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia, Inc.



COSMO BIO Co., LTD.
Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

http://www.cosmobio.co.jp/index_e.asp

E-mail: export@cosmobio.co.jp

Phone : +81-3-5632-9617

FAX : +81-3-5632-9618