

anti-CD40 antibody, mouse monoclonal (5C3)

72-030 100 μ g (US\$ 210)

Shipment and Storage: Ship at 4°C and stored at -20°C. Do not freeze below -20°C

Reactivity: Human

Immunogen: Recombinant extracellular domain of CD40

Applications:

1. Flow cytometry
2. Immunofluorescence staining and Immunocytochemistry (1/100)
3. Immunohistochemistry (acetone or zinc-fixed or frozen section; indirect immuno-staining) (1/100)
4. Stimulation of proliferation of B cell and dendritic cells (20 μ g/ml). Enhancement of proinflammatory cytokine production in human monocytes, such as TNF- α , IL-6 and IL-8

Isotype: Mouse IgG1 κ

Product: Purified from hybridoma culture grown in serum-free medium by salting-out and ion-exchange chromatography.

Form: 1mg/ml in PBS, 50% glycerol, filter-sterilized. Azide- and carrier-free

Background: CD40 is a 45-50-kDa glycoprotein belonging to the tumor necrosis factor (TNF) receptor superfamily. **CD40** is specifically expressed on the surface of B cells and specialized antigen-presenting cells such as dendritic cells and macrophages. **CD40** interacts with the CD40 ligand (CD154) which is found primarily on T cells, playing a role in both humoral and cell-mediated immune responses. Activation of **CD40** on B cells by CD40 ligand causes B cell proliferation, differentiation, immunoglobulin isotype switching, germinal center formation, and stimulation of the humoral memory response. **CD40** has been found to mediate a broad variety of immune and inflammatory responses. Within the cell, the **CD40** molecule acts as a transmembrane signal transducer that leads to activation of intracellular kinases and transcription factors.

Data Link: Swiss-Prot [P25942](#)

References: This antibody is used in ref.2 and 3.

1. Inui S *et al* (1990) "Identification of the intracytoplasmic region essential for signal transduction through a B cell activation molecule, CD40." *Eur J Immunol* **20**: 1747-1753 PMID: [1698631](#) **FC, Stimulation of B cell proliferation.**
2. Yasui T *et al* (2002) "Dissection of B cell differentiation during primary immune responses in mice with altered CD40 signals." *Int Immunol* **14**: 319-329 PMID: [11867568](#) **FC**
3. Ishida I *et al* (2003) "Involvement of CD100, a lymphocyte semaphoring, in the

activation of the human immune system via CD72: implications for the regulation of immune and inflammatory responses.” *Int Immunol.* **15**: 1027-1034 PMID: [12882840](#) FC, Stimulation of B cell proliferation.




Fig.1 Expression of CD40 on the cell surface of monocyte derived dendric cells as analyzed by Flow Cytometry with anti-CD40 antibody (5C3).

Monocyte-derived dendric cells from healthy adult donor were stained with anti-hCD40 antibody (solid line) or with isotype control (broken line).




Fig.2 Immunofluorescent staining of CD40 in MCF7 cells with anti-CD40 antibody (5C3)

Cells were fixed in 4% paraformaldehyde and permeabilized in 0.25% Triton X-100.

Anti-CD40 was used at 1/100 dilution and as 2nd antibody, Alex 488 conjugated goat anti-mouse IgG was used at 1/1,000 dilution. DNA was stained with DAPI.

Related Products: #72-031 anti-CD40 antibody (5C3), Biotin-conjugated.

#72-031 anti-CD40 antibody (5C3), FITC-conjugated.