



Anti Dysbindin-1

Background :

Dysbindin (UniProt: Q96EV8; also known as Biogenesis of lysosome-related organelles complex 1 subunit 8, BLOC-1 subunit 8, Dysbindin-1, Dystrobrevin-binding protein 1, Hermansky-Pudlak syndrome 7 protein, HPS7 protein) is encoded by the DTNBP1 (also known as BLOC1S8, My031) gene (Gene ID: 84062) in human. Dysbindin is a component of the BLOC-1 complex, a complex that is required for normal biogenesis of lysosome-related organelles (LRO), such as platelet dense granules and melanosomes. In concert with the AP-3 complex, the BLOC-1 complex is required to target membrane protein cargos into vesicles assembled at cell bodies for delivery into neurites and nerve terminals. The BLOC-1 complex, in association with SNARE proteins, is also proposed to be involved in neurite extension. It plays a role in synaptic vesicle trafficking and in neurotransmitter release. 3 isoforms produced by alternative splicing and alternative initiation have been described. Isoform 1 is mainly cytoplasmic, but shuttles between the cytoplasm and nucleus. It is exported out of the nucleus via its nuclear export sequence (NES). Its nuclear localization is required for regulation of the expression of genes, such as SYN1. It is detected in neuron cell bodies, axons and dendrites and is mainly located to the postsynaptic density. Isoform 2 also shuttles between the cytoplasm and nucleus and is mainly expressed in the dendritic spine. It is predominantly a synaptic vesicle isoform but also highly expressed in the nucleus. Isoform 3 is exclusively cytoplasmic and is predominantly found in the postsynaptic density (PSD) with little association with synaptic vesicles. Mutations in DTNBP1 gene have been linked to Hermansky-Pudlak syndrome 7 (HPS7) that is characterized by oculocutaneous albinism, bleeding due to platelet storage pool deficiency, and lysosomal storage defects. Defects in DTNBP1 are associated with susceptibility to schizophrenia.

Applications:	Western Blotting (WB)	: 0.3 ug/ml
	Immunofluorescence (IF)	: 1 ug/ml
	Immunohistochemistry (Paraffin) (IHC (P))	: 1 ug/ml
Specificity:	Dysbindin-1	
Immunogen:	MBP-tagged full length recombinant human Dysbindin.	
Host:	Rabbit	
Reactivity:	Human, Mouse, Rat	
Clonality:	Polyclonal	
Subclass:	IgG	
Purification method:	Affinity purified	
Form:	Liquid (0.1M NaPB, pH7.0, 20mg/ml BSA, 0.1% Sodium Azide (NaN ₃) added)	
Conjugation:	None	
Volume:	50 ug (1 mg/ml)	
Storage condition:	-20°C	

Example Assay Data:

1. Immunohistochemistry for formalin/PFA-fixed paraffin embedded sections




Figure 1. IHC(P) result of Anti Dysbindin-1 antibody

2. Western Blotting




Figure 2. Western Blotting results of Anti Dysbindin-1 antibody

References:

Ito H, Morishita R, Shinoda T, Iwamoto I, Sudo K, Okamoto K, Nagata KI. (2010) Dysbindin-1, WAVE2 and Abi-1 form a complex that regulates dendritic spine formation. Mol Psychiatry 15:976-986.

Ito H, Morishita R, Nagata K. (2016) Schizophrenia susceptibility gene product dysbindin-1 regulates the homeostasis of cyclin D1. Biochim Biophys Acta. 1862: 1383-1391.

For research use only, Not for diagnostic use.



COSMO BIO Co., LTD.

2-20, Toyo 2-Chome, Koto-ku, Tokyo 135-0016, JAPAN

T: +81-3-5632-9617 日本のお客様 T: 03-5632-9610

F: +81-3-5632-9618 F: 03-5632-9619

E: export@cosmobio.co.jp W: www.cosmobio.com W: www.cosmobio.co.jp