



MONOCLONAL ANTIBODY

For research use only. Not for clinical diagnosis

Catalog No. CEC-003

Anti- Histone H3 T32ph

BACKGROUND


Post-translation modifications of histones modulate the accessibility and transcriptional competence of specific chromatin regions within the eukaryotic genome. Phosphorylation of histone H3 is unique in the sense that it associates on one hand with open chromatin during gene activation and marks on the other hand highly condensed chromatin during mitosis.

Product type	Primary antibody
Immunogen	Synthetic peptide corresponding to N-terminus region Thr32ph (aa 21-39) of human Histone H3, ATKAARKSAPS(phT)GGVKKPH
Host	Rat
Clone number	6C7G12
Isotype	IgG2a, κ
Source	Culture supernatant
Purification	Ion-exchange chromatography
Form	Liquid
Presentation	Purified monoclonal antibody in PBS, 50% Glycerol, 0.05%w/v ProClin300
Concentration	1 mg/mL
Volume	100 μ L
Label	Unlabeled
Specificity	Histone H3 T32ph Epitope : phosphorylated Thr32 of Histone H3
Cross reactivity	Human, Monkey, Mouse, Rat, Hamster Other species have not been tested.
Storage	Store below -20°C (below -70°C for prolonged storage) Aliquot to avoid cycles of freeze/thaw.
Other	Data Link: UniProtKB/Swiss-Prot P68431 * recommended positive controls is mammalian cell

Application notes	Recommended use WB, ICC Not tested for other applications. Recommended dilutions Western blotting, 1/500 (Fig.2) Immunocytochemistry, 1/500 (Fig.3) Optimal dilutions/concentrations should be determined by the end user.
--------------------------	---

References	1) Strahl and Allis, Nature. 2000 Jan 6;403(6765):41-5. PMID: 10638745
-------------------	--

ANTIBODY CHARACTERIZATION



H3.1	21 ATKAARKS	APAT	GGVKKPH	39
H3.1 S28ph	21 ATKAARKS	phAPAT	GGVKKPH	39
H3.1 T32ph	21 ATKAARKS	APAT	phGGVKKPH	39
H3.3	21 ATKAARKS	APST	GGVKKPH	39
H3.3 S28ph	21 ATKAARKS	phAPST	GGVKKPH	39
H3.3 T32ph	21 ATKAARKS	APST	phGGVKKPH	39
H3 T11ph	1	ARTKQTARKST	phGGKAPRKQC	19

Fig.1 The composition of Histone H3 peptides and the reactivity of Histone H3 T32ph antibody, 6C7G12.




Fig.2 Western blot analysis of the treated-cell extracts using Histone H3 T32ph antibody, 6C7G12




Fig.3 Immunocytochemical analysis of HeLa Cell using Histone H3 T32ph antibody, 6C7G12. (left : interphase, right : metaphase)

For research use only, Not for diagnostic use.



COSMO BIO Co., LTD.

[JAPAN]
TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME,
KOTO-KU. TOKYO 135-0016, JAPAN
Phone: +81-3-5632-9610
FAX: +81-3-5632-9619
URL: <https://www.cosmobio.co.jp/>



COSMO BIO USA

[Outside Japan]
2792 Loker Ave West, Suite 101
Carlsbad, CA 92010, USA
email: info@cosmobioussa.com
Phone/FAX: (+1) 760-431-4600
URL: www.cosmobioussa.com