



MONOCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. CTB-LC3-2-IC

Anti LC3 (Clone: LC3·1703)

BACKGROUND

LC3B is one of the mammalian Atg8 homologs and widely used as an autophagosome marker. Immediately after synthesis, LC3 is processed by Atg4 and becomes LC3-I. Upon induction of autophagy, the C-terminal glycine of LC3-I is conjugated to phosphatidylethanolamine, resulting in formation of membrane-bound LC3-II. Most LC3-II is thought to be present on autophagosome membrane. The autophagosome subsequently fuses with a lysosome, where inside materials, including LC3-II, are degraded. The expression level of LC3-II generally correlates with the number of autophagosome.

Product type	Primary antibodies
Host	Mouse
Source	
Form	Liquid Protein G purified PBS (pH7.4) with 1% BSA and less than 0.1% NaCl as a preservative.
Volume	500 μ l
Concentration	0.1mg/ml
Specificity	LC3
Antigen	Human Recombinant LC3
Clone	LC3·1703
Isotype	IgG1

Application notes ICC, Immuno EM
Recommended use

Recommended dilutions

ICC: 1/100
Immuno EM : 1/10

Optimal dilutions/concentrations should be determined by the end user.

Staining Pattern


Cross reactivity Human, Mouse

Storage Store below -20° C (below -70° C for prolonged storage).

Aliquot to avoid cycles of freeze/thaw.


References

- 1) Kabeya, Y., Mizushima, N., Ueno, T., Yamamoto, A., Kirisako, T., Noda, T., Komiyama, E., Ohsumi, Y. and Yoshimori, T. LC3, a mammalian homologue of yeast Apg8p, is localized in autophagosome membranes after processing EMBO J. 19, 5720-5728. (2000)
- 2) Mizushima, N., Yoshimori, T. How to interpret LC3 immunoblotting Autophagy 3:542-545 (2007)
- 3) Mizushima, N., Yoshimori, T. and Levine, B. Methods in mammalian autophagy research. Cell 140; 313-326 (2010)



(Fig 1) Immunofluorescence microscopy analysis of mouse embryonic fibroblasts (MEFs).

MEFs were cultured in regular DMEM supplemented with 10% FBS (left) or DMEM without amino acids (right) for 1 hr. After fixation with 4% paraformaldehyde for 10 min at room temperature, they were permeabilized with 50 μ g/ml digitonin for 5 min. Cells were then subjected to immunofluorescence microscopy using #1703 anti-LC3 antibody at 1:100 dilution. Scale bar, 20 μ m.



(Fig 2) Immuno-electron microscopy analysis of mouse embryonic fibroblasts (MEFs).

MEFs were cultured in DMEM without amino acids for 2 hr. After fixation with 4% paraformaldehyde for 2 hr at room temperature, they were permeabilized with liquid nitrogen. Immuno-electron microscopy analysis (pre-embedding method) of endogenous LC3 was performed using #1703 anti-LC3 antibody at 1:10 dilution. The gold labeling was intensified by using a silver enhancement kit (HQ silver enhancement kit, Nano probes, NY). Scale bar, 500 nm.

For research use only. Not for clinical diagnosis.



COSMO BIO Co., LTD.

【JAPAN】

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME,
KOTO-KU. TOKYO 135-0016, JAPAN
Phone: +81-3-5632-9610
FAX: +81-3-5632-9619
URL: <https://www.cosmobio.co.jp/>



COSMO BIO USA

【Outside Japan】

2792 Loker Ave West, Suite 101
Carlsbad, CA 92010, USA
email: info@cosmobioussa.com
Phone/FAX: (+1) 760-431-4600
URL: www.cosmobioussa.com