

POLYCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. KZ-HS-P08


Anti Bovine Follistatin

BACKGROUND

Follistatin also known as activin-binding protein is a protein that in humans is encoded by the *FST* gene. Follistatin is an autocrine glycoprotein that is expressed in nearly all tissues of higher animals. It was initially isolated from follicular fluid and was identified as a protein fraction that inhibited follicle-stimulating hormone (FSH) secretion from the anterior pituitary, and so was known as FSH-suppressing protein (FSP). Since then its primary function has been determined to be the binding and bioneutralization agent of members of the TGF- β superfamily, with primary focus on activin.

Product type	Primary antibodies
Host	Rabbit
Form	Serum
Volume	50 µl
Antigen	Bovine Follistatin
Cross reactivity	Bovine, Rat, Mouse
Application notes	Recommended dilutions

RIA : 1/2,000 - 1/10,000 IHC : 1/200 - 1/1,000 WB : 1/200 - 1/1,000



Elution profile on reverse-phase HPLC of the partially purified activin-follistatin complex of fraction 2 in figure 6 applied to a Cosmosil 5C₁₈-AR column (10×250 mm). The column was developed with the indicated gradient of acetonitrile (ACN) in 0.1% trifluoroacetic acid.

Storage	Store below -20°C (below -70°C for prolonged storage). Aliquot to avoid cycles of freeze/thaw.
References	 Hasegawa Y, Eto Y, Ibuki Y & Sugino H (1994) Activin as autocrine and paracrine factor in the ovary. Horm Res 41(Suppl 1): 55-62. PMID: <u>8088695</u>.

For research use only. Not for clinical diagnosis.



(JAPAN)

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME, KOTO-KU. TOKYO 135-0016, JAPAN Phone: +81-3-5632-9610 FAX: +81-3-5632-9619 URL: https://www.cosmobio.co.jp/



[Outside Japan] 2792 Loker Ave West, Suite 101 Carlsbad, CA 92010, USA email: info@cosmobiousa.com Phone/FAX: (+1) 760-431-4600 URL: www.cosmobiousa.com