

## **POLYCLONAL ANTIBODY**

For research use only. Not for clinical diagnosis.

#### Catalog No. NIG-L1-SHA1

# Anti-dSSRP1

#### **BACKGROUND**

**dSSRP1** is a *Drosophila* counterpart of the FACT subunit. FACT consists of a heterodimer of **SSRP1** and SPT16 and was identified as a factor that *facilitates chromatin transcription in vitro*. Further studies revealed that the factor also facilitates various chromatin transactions such as transcriptional regulation, DNA replication and histone replacement. The antibodies against **dSSRP1** are powerful tool for Western blots(Ref.1), immunostaining(Ref.2), immunoprecipitation and chromatin immunopreciptation(Ref.1,2,3) in studies using *Drosophila*.

Product type Primary antibodies

Antigen Bacterially expressed N-terminal region of dSSRP1 (amino acid residues 1-420)

Host Rabbit
Clone Isotype Source Serum
Form Liquid

Form Liquid
Concentration Volume 100 ul
Label Unlabeled
Specificity dSSRP1
Cross reactivity Drosophila

**Storage** Store at 4°C for a short time (below -70°C for prolonged storage).

Aliquot to avoid cycles of freeze/thaw.

Other

#### **Application notes**

Recommended dilutions

Western Blotting: 1/4,000 - 1/8,000. Predicted molecular weight: about 93kDa

· Immunohistochemistry: 1/500 - 1/2,000

• ELISA: 1/4,000 - 1/10,000

Immunoprecipitation: 0.5 – 2 ul

Chromain immunoprecipitation: 1 - 4 ul

Other applications have not been tested.

Optimal dilutions/concentrations should be determined by the end user.

#### **Staining Pattern**

The antibodies stain nuclei of embryos, larval, pupal and adult tissues.

In the adult oocytes, a highly phosphorylated inactive form is deposited in the cytoplasm.

## References

 SHIMOJIMA, T. et al. Drosophila FACT contributesto Hox gene expression through physical and functional interactions with GAGA factor. Genes & Dev. 17,1605-1616, 2003

- 2) Saunders, A. et al. Tracking FACT and RNA Polymerase IIElongation Complex Through hChromatin in Vivo. Science, 301, 1094-1096, 200
- 3) Nakayama, T. et al. Drosophila GAGA factor directs histone H3.3 replacement that prevents the heterochromatin spreading. Genes & Dev. 21, 552-561, 2007




Fig.1 Immunostaining of a polytene chromosome(anti dSSRP1 antibody)

## For research use only. Not for clinical diagnosis.



# COSMO BIO CO., LTD.

[JAPAN]

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME, KOTO-KU. TOKYO 135-0016, JAPAN Phone: +81-3-5632-9610 FAX: +81-3-5632-9619

URL: https://www.cosmobio.co.jp/



# COSMO BIO USA

[Outside Japan] 2792 Loker Ave West, Suite 101 Carlsbad, CA 92010, USA email: info@cosmobiousa.com Phone/FAX: (+1) 760-431-4600 URL: www.cosmobiousa.com