

POLYCLONAL ANTIBODY

Catalog No. SDT-01-APX

Anti-Cytosolic Ascorbate Peroxidase

| Product type Host Source Form Volume Concentration Specificity Antigen Isotype | Primary antibodies Rabbit Serum (Partially purified (0-40% ammonium sulfate fraction)) TBS (pH7.5) with 0.02% NaN3 as a preservative. 200ul Cytosolic Ascorbate Peroxidase Cytosolic ascorbate peroxidase purified from Zea mays IgG |
|--|---|
| Application notes | WB, IHC IP, ELISA, FC: not tested Recommended use Recommended dilutions Western Blot: 1/1,000-1/2,000. Predicted molecular weight: 145 kDa 28kDa Immunohistochemistry: 1/100-1/500 |
| | Optimal dilutions/concentrations should be determined by the end user. Staining Pattern |
| Cross reactivity | Plant (Maize, Rice, etc.) |
| Storage | Store below -20°C (below -70°C for prolonged storage). |
| | Aliquot to avoid cycles of freeze/thaw. |
| References | Koshiba, T. (1993) Cytosolic ascorbate peroxidase in seedlings and leaves of maize (Zea mays). Plant Cell Physiol. 34:713-721. |

For research use only. Not for clinical diagnosis.

This product was made by using the research results of Tokyo Metropolitan University.

