

### POLYCLONAL ANTIBODY

For research use only, Not for diagnostic use

## Catalog No. TIP-SN-P05

# Anti α-Synuclein (41-50)

#### **BACKGROUND**

Alpha-Synuclein, a 140-amino acid protein abundantly expressed in presynaptic terminals, is known as a component of intraneuronal or glial inclusions observed in cases of Parkinson's disease (PD), Dementia with Lewy bodies (DLB) and Multiple system atrophy (MSA). Alpha-Synuclein is a natively unfolded protein, however, fibrillization or conformational change of Alpha-synuclein is central in the pathogenesis of Alpha-synucleinopathies. The amino-terminal region of Alpha-synuclein consists of seven imperfect repeats, each 11 amino acids in length, with the consensus sequence KTKEGV. The repeats partially overlap with a hydrophobic region (amino acids 61-95). The carboxy-terminal region (amino acids 96-140) is negatively charged. These antibodies are powerful tools for biochemical and IHC analyses of neurodegenerative diseases and for evaluation of conformational changes of Alpha-synuclein.

Primary antibody **Product type** 

GSKTKEGVVHC (α-syn 41-50) **Immunogen** Raised in Rabbit (New Zealand White)

Source Anti-serum

**Purification** 

**Form** Liquid. Anti-serum with 0.1% NaN3 as a preservative.

Concentration

50 uL Volume Unlabeled Label **Specificity** α-Syn

**Cross reactivity** Human, Mouse

Store below -20°C. (below -70°C for prolonged storage). Storage

Aliquot to avoid cycles of freeze/thaw.

**Application notes** •ELISA: 1/500-1/2000

Recommended Western Blot: 1/500-1/1000

dilutions •Immunohistochemistry:1/500-1/2000

Other applications have not been tested. Optimal dilutions/ concentrations

should be determined by the end user.

References 1)Masami Masuda, et al. Inhibition of a-synuclein fibril assembly by small molecules:

Analysis using epitope-specific antibodies, FEBS Letters (2009) 583, 787-791, PMID

19183551



Cosmo Bio Co., Ltd.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: http://www.cosmobio.co.jp e-mail: export@cosmobio.co.jp

[Outside Japan] Phone: +81-3-5632-9617 [国内連絡先] Phone: +81-3-5632-9610 FAX: +81-3-5632-9618 FAX: +81-3-5632-9619

# **RELATED PRODUCTS:**

| Product Name                        | Quantity | Maker | Cat#       |
|-------------------------------------|----------|-------|------------|
| Anti α-Synuclein (1-10)             | 50 uL    | CAC   | TIP-SN-P01 |
| Anti α-Synuclein (11-20)            | 50 uL    | CAC   | TIP-SN-P02 |
| Anti α-Synuclein (21-30)            | 50 uL    | CAC   | TIP-SN-P03 |
| Anti α-Synuclein (31-40)            | 50 uL    | CAC   | TIP-SN-P04 |
| Anti α-Synuclein (51-60)            | 50 uL    | CAC   | TIP-SN-P06 |
| Anti α-Synuclein (61-70)            | 50 uL    | CAC   | TIP-SN-P07 |
| Anti α-Synuclein (75-91)            | 50 uL    | CAC   | TIP-SN-P08 |
| Anti α-Synuclein (131-140)          | 50 uL    | CAC   | TIP-SN-P09 |
| Anti α-Synuclein (9 Antibodies Set) | 9*10 uL  | CAC   | TIP-SN-SET |

For research use only, Not for diagnostic use.



# COSMO BIO CO., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN e-mail: export@cosmobio.co.jp

URL: http://www.cosmobio.co.jp [Outside Japan] Phone : +81-3-5632-9617 [国内連絡先] Phone: +81-3-5632-9610 FAX: +81-3-5632-9618 FAX: +81-3-5632-9619