

Anti human ER alpha mouse monoclonal antibody

ER alpha: Estorgen Receptor alpha

Code No	PP-H4624-00	Application / Recommended Concentration In order to obtain the best results, optimal work determined by each individual user.	
Clone No.	H4624	Western Blot 1 ug/mL	
Lot.	***	Non-reducing Western Plet	
Concentration	1 mg/mL	Non reducing Western Blot Not yet tested	
Volume	100 uL	ELISA 0.3 ug/mL (A ²	
lg Class	G2a	Immunoprecipitation Decide by use	
, , , , , , , , , , , , , , , , , , ,	Estrogen receptor alpha (ERa, ER; NR3A1) is a member of steroid receptor (AR, GR, MR, PR). The natural ligand for ER is the classical estrogenic	Supershift Assay Not yet tested	
	compound 17b-estradiol. ERa is expressed in a wide variety of tissues. Expression was found at variable levels in bones, in uterus, bladder, ovary, prostate,	Chromatin immunoprecipitation Not yet tested	
	testis, epididymys, kidney, breast, heart, vessel wall, pituitary and hypothalamus. ER has a variety of central physiological roles, including those involved in	Immunohistochemistry 10-20 ug/mL	
	maintenance of the reproductive, cardiovascular, musculoskeletal and central nervous systems. ERa has		

Nomenclature	NR3A1
Genbank	M12674
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human ER alpha (2-180 aa) .
Specificity	This antibody specifically recognizes humman ER alpha and cross reacts with mouse and rat ER alpha. This antibody does not recognize human ER beta.

been shown to form homodimers as well as heterodimers with ERb. Both ERa and ERb can give


rise to numerous isoforms.


preservative.


Ammonium sulfate fractionation

the best results, optimal working dilutions should be ach individual user.

Western Blot	1 ug/mL
Non reducing Western Blot	Not yet tested
ELISA	0.3 ug/mL (A450=0.3)
Immunoprecipitation	Decide by use
Supershift Assay	Not yet tested







Human Uterus Endometrial epithelium cell paraffin section

Storage

Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Reference

Notes Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited. MADE IN JAPAN

Apr 17, 2008



Purification

Formulation

Distributor:

Physiological saline with 0.1% NaN3 as a

2-20, Toyo, 2Chome, Tokyo 135-0016, Japan TEL: +81-3-5632-9617 FAX: +81-3-5632-9618 http://www.cosmobio.co.jp



Manufactured by Perseus Proteomics Inc. 4-7-6, Komaba, Meguro-ku, Tokyo 153-0041, Japan http://www.ppmx.com