



Anti Galanin (Porcine) Serum

Cat. No. YII-Y180-EX Lot No. 198571025

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic galanin (porcine) peptide. The product vial contains 50 µL of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with galanin (porcine).

Immunogen: Synthetic galanin (porcine), carrier free **Host:** Rabbit

Amino Acid Sequence of Galanin (porcine)¹⁾ : GWTLNSAGYL LGPHAIDNHR
SFHDKYGLA-NH₂

Product Form: Lyophilized unpurified serum **Size:** 50 µL

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:2,500 (final dilution ~1:17,500) for radioimmunoassay; 1:500-2,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): Galanin (porcine) 100%, galanin (3-29) (Porcine) 100%, galanin (10-29) (porcine) 100%, galanin (15-29) (porcine) 100%, GRP (porcine) 0%, neurokinin A 0%, neurokinin B 0%, neuropeptide B 0%, substance P 0%, PP (human) 0%, glucagon 0%, VIP (porcine) 0%.

Positive Control (immunohistochemistry): Rat duodenum

Species Tested: Rat²⁾, tuna fish

REFERENCES:

- 1) K. Tatemoto, A. Rokaeus et al., Galanin-a novel biologically active peptide from porcine intestine. FEBS Letters 164: 124-128, 1983
- 2) Y. Takeda, N. Yanaihara et al., Inhibitory effect of novel neuropeptide, galanin on glucose-induced insulin release in isolated perfused rat pancreas. Proceedings of 8th Gut Hormone Conference, Japan Society of Gut Hormones (Ed.) 6: 427-433, 1986

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

