



Anti Preprorenin (21-64) (Human) Serum Cat. No. YII-Y190-EX Lot No. 006771016

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic preprorenin (21-64) (human) peptide. The product vial contains 50µL of the titled antiserum obtained by lyophilizing its 0.001M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with preprorenin (human) and prorenin (human).

Immunogen: Synthetic preprorenin (21-64) (human), carrier free Host: Rabbit

Amino Acid Sequence of Preprorenin (21-64) (human)1), 2):

21 64

TFGLPTDTTT FKRIFLKRMP SIRESLKERG VDMARLGPEW SQPM

Product Form: Lyophilized unpurified serum Size: 50 μL

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN3 0.1%).

Storage: The product will be stable for over one year if stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Reconstituted antiserum solution can also be aliquotted and stored at -20°C to -80°C for six months without marked loss of activity. Repeated freezing- thawing should be avoided.

Suggested Working Dilution Range: 1:1,000 ~ 1:5,000 for enzyme immunoassay; 1:200 ~ 1:2,000 for immunohistochemistry (frozen section). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on non-competitive EIA): preprorenin (21-64) (human) 100%, prorenin (265-294) (human) 0%, prorenin (314-337) (human) 0%, prorenin (human, recombinant) 100%, renin (human, recombinant) 0%

Positive Control (immunohistochemistry): Human kidney 2)

Species Tested: Human

REFERENCES:

- 1) T. Imai, H. Miyazaki, et al., Cloning and sequence analysis of cDNA for human renin precursor. Proceedings of National Academy of Sciences, USA 80: 7405-7409, 1983
- 2) C. Yanaihara, M. Kadowaki et al., Production of region-specific antisera to human prorenin and renin using synthetic peptides. Biomedical Research 8: 95-101, 1987

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

