



Anti Orexin B (Mouse, Rat) Serum Cat. No. YII-Y451-EX Lot No. 1385170126

Description: This antiserum was raised in a rabbit by immunization with a keyhole limpet hemocyanin (KLH) conjugate of synthetic orexin B (mouse, rat) peptide. The product vial contains 50mL of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5 mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with orexin B (mouse, rat).

Immunogen: Synthetic Orexin B (mouse, rat)-KLH conjugate

Host:Rabbit

Amino Acid Sequence of Orexin B (mouse, rat)1) 2):

RPGPPGLQGR LQRLLQANGN HAAGILTM-NH2

Product Form: Lyophilized unpurified serum Size: 50mL

Reconstitution: Reconstitute the product with 0.5 mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at $-20\Box$ to $-80\Box$ until opened. Upon reconstitution, the antiserum solution must be stored at $2\Box$ to $8\Box$ and used within one month.Repeated freezing thawing should be avoided.

Suggested Working Dilution Range: 1:8,000 for enzyme immunoassay (EIA); 1:6,000 for immunohistochemistry. Optimal dilution should be determined by each laboratory for each application.

Specificity(based on EIA): Orexin B (mouse, rat) 100%, Orexin A (bovine, dog, human, mouse, porcine, rat) 0%

Positive Control (Immunohistochemistry): Hypothalamus

Species Tested (Immunohistochemistry): Mouse

Related Peptide: Orexin A (bovine, dog, human, mouse, porcine, rat):

Related Antiserum: Anti Orexin A Serum Y450

REFERENCES:

- 1. T. Sakurai, A. Amemiya et al., Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*, **92**: 573-585, 1998
- 2. L. Lecea, TS. Kilduff et al., The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc. Natl. Acad. Sci. U.S.A.*, **95**:322-327, 1998

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

