








## Recombinant Rhesus Macaque Interleukin-3 protein (IL3) (Active)

Product Code       CSB-AP001981MOW         Abbreviation       Recombinant Rhesus macaque IL3 protein (Active)         Uniprot No.       P25140         Form       Lyophilized powder         Storage Buffer       Lyophilized from a 0.2 μm filtered PBS, pH 7.4, 5 % trehalose         Product Type       Interleukin         Immunogen Species       Macaca mulatta (Rhesus macaque)         Biological Activity       Assay #1: Fully biologically active when compared to standard. The ED50 determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10 <sup>5</sup> IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10 <sup>7</sup> IU/mg.         Purity       >98% as determined by SDS-PAGE.         Sequence       APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
Uniprot No.         P25140           Form         Lyophilized powder           Storage Buffer         Lyophilized from a 0.2 μm filtered PBS, pH 7.4, 5 % trehalose           Product Type         Interleukin           Immunogen Species         Macaca mulatta (Rhesus macaque)           Biological Activity         Assay #1: Fully biologically active when compared to standard. The ED50 determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10⁵ IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10⁻ IU/mg.           Purity         >98% as determined by SDS-PAGE.           Sequence         APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
Form         Lyophilized powder           Storage Buffer         Lyophilized from a 0.2 μm filtered PBS, pH 7.4, 5 % trehalose           Product Type         Interleukin           Immunogen Species         Macaca mulatta (Rhesus macaque)           Biological Activity         Assay #1: Fully biologically active when compared to standard. The ED50 determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10 <sup>5</sup> IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10 <sup>7</sup> IU/mg.           Purity         >98% as determined by SDS-PAGE.           Sequence         APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
Storage Buffer         Lyophilized from a 0.2 μm filtered PBS, pH 7.4, 5 % trehalose           Product Type         Interleukin           Immunogen Species         Macaca mulatta (Rhesus macaque)           Biological Activity         Assay #1: Fully biologically active when compared to standard. The ED50 determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10 <sup>5</sup> IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10 <sup>7</sup> IU/mg.           Purity         >98% as determined by SDS-PAGE.           Sequence         APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
Interleukin   Immunogen Species   Macaca mulatta (Rhesus macaque)	ssay
Immunogen Species  Macaca mulatta (Rhesus macaque)  Assay #1: Fully biologically active when compared to standard. The ED50 determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10 <sup>5</sup> IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10 <sup>7</sup> IU/mg.  Purity  >98% as determined by SDS-PAGE.  Sequence  APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
Assay #1: Fully biologically active when compared to standard. The ED50 determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10 <sup>5</sup> IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10 <sup>7</sup> IU/mg.  Purity >98% as determined by SDS-PAGE.  Sequence APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
determined by the dose- dependent stimulation of the proliferation of murin NFS-60 cells is less than 5.0 ng/ml, corresponding to a specific activity of >2x10 <sup>5</sup> IU/mg. Assay #2: The ED50 as determined by a cell proliferation as using human TF-1 cells is less than 0.05 ng/ml, corresponding to a specific activity of >2.0x10 <sup>7</sup> IU/mg.  Purity >98% as determined by SDS-PAGE.  Sequence APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	ssay
Sequence  APMTQTTSLK TSWAKCSNMI DEIITHLNQP PLPSPDFNNL NEEDQTILV KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	
KNLRRSNLEA FSKAVKSLQN ASAIESILKN LPPCLPMATA APTRPPIRIT NGDRNDFRRK LKFYLKTLEN EQAQ	
	E
Research Area Immunology	
Source E.coli	
Target Names IL3	
Expression Region 20-143aa	
Notes Repeated freezing and thawing is not recommended. Store working aliquo 4°C for up to one week.	s at
Tag Info Tag-Free	
Mol. Weight 14.0 kDa	
Protein Length Full Length of Mature Protein	
<b>PubMed ID</b> 2251147; 2257297	
Image	









## **Endotoxin**

Less than 1.0 EU/µg as determined by LAL method.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

## Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.