





# Recombinant Human ATP-sensitive inward rectifier potassium channel 10 (KCNJ10), partial


<b>Product Code</b>	CSB-EP012048HU1
<b>Relevance</b>	May be responsible for potassium buffering action of glial cells in the brain. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of Extracellular domain potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. Can be blocked by Extracellular domain barium and cesium .
<b>Abbreviation</b>	Recombinant Human KCNJ10 protein, partial
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	P78508
<b>Alias</b>	ATP-dependent inwardly rectifying potassium channel Kir4.1Inward rectifier K(+) channel Kir1.2;Potassium channel, inwardly rectifying subfamily J member 10
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	Greater than 90% as determined by SDS-PAGE.
<b>Sequence</b>	FLAKIARPKKRAETIRFSQHAVVASHNGKPCLMIRVANMRKSLIGCQVTGKLL QTHQTKEGENIRLNQVNVTFQVDTASDSPFLILPLTFYHVVDTSPLKDLPLRS GEGDFELVLILSGTVESTSATCQVRTSYLPEEILWGYEFTPAISLSASGKYIADF SLFDQVVKVASPSGLRDSTVRYGDPEKLKLEESLREQAEKEGSALSVRISNV
<b>Research Area</b>	Transport
<b>Source</b>	E.coli
<b>Target Names</b>	KCNJ10
<b>Expression Region</b>	165-379aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	N-terminal 6xHis-SUMO-tagged
<b>Mol. Weight</b>	39.8kDa
<b>Protein Length</b>	Cytoplasmic Domain
<b>Image</b>	



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.



Based on the SEQUEST from database of E.coli host and target protein, the LC-MS/MS Analysis result of CSB-EP012048HU could indicate that this peptide derived from E.coli-expressed Homo sapiens (Human) KCNJ10.



Based on the SEQUEST from database of E.coli host and target protein, the LC-MS/MS Analysis result of CSB-EP012048HU could indicate that this peptide derived from E.coli-expressed Homo sapiens (Human) KCNJ10.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

## Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.