




# Recombinant Human Renin receptor (ATP6AP2)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-YP002384HU  |
| <b>Relevance</b>         | Functions as a renin and prorenin cellular receptor. May mediate renin-dependent cellular responses by activating ERK1 and ERK2. By increasing the catalytic efficiency of renin in AGT/angiotensinogen conversion to angiotensin I, it may also play a role in the renin-angiotensin system (RAS).   |
| <b>Abbreviation</b>      | Recombinant Human ATP6AP2 protein   |
| <b>Storage</b>           | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.   |
| <b>Uniprot No.</b>       | O75787  |
| <b>Alias</b>             | ATPase H(+)-transporting lysosomal accessory protein 2 ATPase H(+)-transporting lysosomal-interacting protein 2 ER-localized type I transmembrane adaptor Embryonic liver differentiation factor 10 N14F Renin/prorenin receptor Vacuolar ATP synthase membrane sector-associated protein M8-9  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Homo sapiens (Human)  |
| <b>Purity</b>            | Greater than 90% as determined by SDS-PAGE.   |
| <b>Sequence</b>          | NEFSILKSPGSSVFRNGNWPIGERIPDVAALSMGFSVKEDLSWPGLAVGNLF<br>HRPRATVMVMVKGVNKLALPPGSVISYPLENAPFSLDSVANSIHSLSFSEETPV<br>VLQLAPSEERVYMGKANSVFEDLSVTLRQLRNRLFQENSVLSSLPLNLSLRN<br>NEVDLLFLSELQVLHDISSLLSRHKHLAKDHSPDLYSLELAGLDEIGKRYGEDS<br>EQFRDASKILVDALQKFADDMYSLYGGNAVVELVTVKSFDTSIRKTRTILEAK<br>QAKNPASPYNLAYKYNFEYSVVFNMVLWIMIALALAVIITSYNIWNMDPGYDSII<br>YRMTNQKIRMD |
| <b>Research Area</b>     | Signal Transduction   |
| <b>Source</b>            | Yeast   |
| <b>Target Names</b>      | ATP6AP2   |
| <b>Protein Names</b>     | Recommended name: Renin receptor<br>Alternative name(s): ATPase H(+)-transporting lysosomal accessory protein 2 ATPase H(+)-transporting lysosomal-interacting protein 2 ER-localized type I transmembrane adaptor Embryonic liver differentiation f  |
| <b>Expression Region</b> | 17-350  |
| <b>Notes</b>             | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| <b>Tag Info</b>          | N-terminal 10xHis-tagged  |
| <b>Mol. Weight</b>       | 40.0kDa   |
| <b>Protein Length</b>    | Full Length of Mature Protein   |



## Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

## Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.