





Recombinant Human Probable global transcription activator SNF2L2 (SMARCA2), partial

Product Code	CSB-YP021799HU
Relevance	Transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. Belongs to the neural progenitors-specific chromatin rodeling complex (npBAF complex) and the neuron-specific chromatin rodeling complex (nBAF complex). During neural development a switch from a st/progenitor to a postmitotic chromatin rodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural st/progenitor cells to postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural st cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth .
Abbreviation	Recombinant Human SMARCA2 protein, partial
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P51531
Alias	ATP-dependent helicase SMAR;CA2BRG1-associated factor 190B;BAF190BProtein brahma homolog
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	SYYTVAHAISERVEKQSALLINGTLKHYQLQGLEWMVSLYNNNLNGILADEMG LGKTIQTIALITYLMEHKRLNGPYLIIVPLSTLSNWTYEFDKWAPSVVKISYKGTP AMRRSLVPQLRSGKFNVLLTTYEYIIKDKHILAKIRWKYMIVDEGHRMKNHHCK LTQVLNTHYVAPRRILLTGTPLQNKLPELWALLNFLLPTIFKSCSTFEQWFNAPF AMTGERVDLNEEETILIIRRLHKVLRPFLLRRLKKEVESQLPEKVEYVIKCDMSA LQKILYRHMQAKGILLTDGSEKDKKGKGGAKTLMNTIMQLRKICNHPYMFQHIE ESFAEHLGYSNGVINGAELYRASGKFELLDRILPKLRATNHRVLLFCQMTSLMT IMEDYFAFRNFLYLRLDGTTKSEDRAALLKKFNEPGSQYFIFLLSTRAGGLGLN LQAADTVVIFDSDWNPHQDLQAQDRAHRIGQQNEVRVLRLCTVNSVEEKILAA AKYKLNVDQKVIQAGMFDQKSSSHERRAF
Research Area	Neuroscience
Source	Yeast








🕜 Tel: +1-301-363-4651 💢 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🌘





Target Names	SMARCA2
Protein Names	Recommended name: Probable global transcription activator SNF2L2 EC= 3.6.4Alternative name(s): ATP-dependent helicase SMARCA2 BRG1-associated factor 190B Short name= BAF190B Protein brahma homolog Short name= hBRM SNF2-a
Expression Region	700-1216aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	61.7kDa
Protein Length	Partial
Image	

Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Shelf Life

storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.

The shelf life is related to many factors, storage state, buffer ingredients,