




# Recombinant Human T cell receptor alpha chain constant (TRAC)

<b>Product Code</b>	CSB-YP024144HU
<b>Abbreviation</b>	Recombinant Human TRAC protein
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	P01848
<b>Product Type</b>	Recombinant Proteins
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	Greater than 90% as determined by SDS-PAGE.
<b>Sequence</b>	PNIQNPDPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKTVLDM RSMDFKSNSAVAWSNKSDFACANAFNNSIIPEDTFFPSPPESSCDVKLVEKSFE TDTNLFQNLQSVIGFRILLKLVAGFNLLMTLRLWSS
<b>Research Area</b>	Immunology
<b>Source</b>	Yeast
<b>Target Names</b>	TRAC
<b>Protein Names</b>	Recommended name: T-cell receptor alpha chain C region
<b>Expression Region</b>	1-142aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	N-terminal 6xHis-tagged
<b>Mol. Weight</b>	17.9kDa
<b>Protein Length</b>	Full Length

## Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.



## Description

Intact human T cell receptor alpha constant (TRAC) cDNA (1-142aa) with an N-terminal 6xHis-tag was expressed in the yeast. The forming protein is the recombinant full-length human TRAC protein. The purity of this protein is greater than 90% determined by SDS-PAGE. Under reducing conditions, the gel showed a molecular weight band of about 16 kDa. This recombinant TRAC protein can be used as the immunogen for antibody synthesis. Besides, it also may be applied in the studies of TRAC-related immunology.

The extracellular region of  $\alpha\beta$  T cell receptors (TCRs) is made up of an  $\alpha$  chain and a  $\beta$  chain, each containing Ig-like variable and constant domains. TRAC, the TCR C $\alpha$  domain, is significant in differing from the classical IgC1 domain. Substantial evidence indicated that TRAC interfaces with CD3. Mutagenesis has demonstrated that the interactions of CD3 $\delta$ ? and CD3 $\gamma$ ? subunits with the TcR C $\alpha$  and C $\beta$  domains, respectively, contribute to the stability and function of the TCR-CD3 signaling complex. Gijs I. van Boxel etc. concluded that the TCR C $\alpha$  domain can also adopt two very different stable conformations, a fundamental property that is potentially central to TCR function and that the TCR C $\alpha$  domain may facilitate the " $\beta$ -strand slippage."

## Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.