





## Recombinant Calloselasma rhodostoma Snaclec rhodocytin subunit beta

| Product Code         | CSB-YP872766CBG   |
|----------------------|---|
| Relevance            | Elicits platelet aggregation by the binding to the C-type lectin domain family 1 member B (CLEC1B/CLEC2). Binding leads to tyrosine phosphorylation in the Cytoplasmic domain tail of CLEC1B, which promotes the binding of spleen tyrosine kinase (Syk), subsequent activation of PLCgamma2, and platelet activation and aggregation. Binding to GPIbalpha (GP1BA) and alpha2/beta-1 (ITGA2/ITGB1) may also induce aggregation, but this is controversial. |
| Abbreviation         | Recombinant Calloselasma rhodostoma Snaclec rhodocytin subunit beta protein   |
| Storage              | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.   |
| Uniprot No.          | Q9I840  |
| Alias                | Aggretin beta chain Rhodoaggretin subunit beta  |
| <b>Product Type</b>  | Recombinant Protein   |
| Immunogen Species    | Calloselasma rhodostoma (Malayan pit viper) (Agkistrodon rhodostoma)  |
| Purity               | Greater than 90% as determined by SDS-PAGE.   |
| Sequence             | DCPSGWSSYEGHCYKPFNEPKNWADAERFCKLQPKHSHLVSFQSAEEADFV<br>VKLTRPRLKANLVWMGLSNIWHGCNWQWSDGARLNYKDWQEQSECLAFRG<br>VHTEWLNMDCSSTCSFVCKFKA   |
| Research Area        | Others  |
| Source               | Yeast   |
| <b>Protein Names</b> | Recommended name: Rhodocytin subunit beta Alternative name(s): Aggretin beta chain  |
| Expression Region    | 24-146aa  |
| Notes                | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.   |
| Tag Info             | N-terminal 6xHis-tagged   |
| Mol. Weight          | 16.4kDa   |
| Protein Length       | Full Length of Mature Protein   |
| Image                |   |




## **CUSABIO TECHNOLOGY LLC**

Tel: +1-301-363-4651 

☐ Email: cusabio@cusabio.com ☐ Website: www.cusabio.com ☐







(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

## **Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.