

Ubiquitin-Lys-TAMRA (5-TAMRA-Lys(Ub)-Gly-OH)

Ubiquitin substrate



Cat. No. 60-0118-050

Lot. No. 30114

Quantity: 50 µg

Storage: -70°C

FOR RESEARCH USE ONLY

NOT FOR USE IN HUMANS

CERTIFICATE OF ANALYSIS Page 1 of 1

Background

A shift in both the excitation and the emission toward longer wavelength helps overcome problems of compound autofluorescence in screening assays. One such substrate is the rhodamine based ubiquitin-εN-(αN-tetramethyl-rhodamine)-lysine (Ubiquitin-Lys-TAMRA), which mimics the naturally occurring isopeptide bond between the C-terminus of ubiquitin and the ε-amino group of a lysine residue of an ubiquitinated protein (Tirat *et al.*, 2005). Cleavage of the isopeptide bond results in a decrease of fluorescence polarization, which makes Ubiquitin-Lys-TAMRA suitable for high-throughput screening applications (Hassiepen *et al.*, 2007). Fluorescence polarization, in contrast to fluorescence intensity, allows a ratiometric read-out of the activity and is thus less sensitive to autofluorescing or quenching caused by test compounds (Tirat *et al.*, 2005).

References:

Hassiepen U, Eidhoff U, Meder G, Bulber JF, Hein A, Bodendorf U, *et al.* (2007) A sensitive fluorescence intensity assay for deubiquitinating proteases using ubiquitin-rhodamine110-glycine as substrate. *Anal Biochem* 371, 201-207.

Tirat A, Schilb A, Riou V, Leder L, Gerhartz B, Zimmermann J, *et al.* (2005) Synthesis and characterization of fluorescent ubiquitin derivatives as highly sensitive substrates for the deubiquitinating enzymes UCH-L3 and USP-2. *Anal Biochem* 343, 244-255.

Physical Characteristics

Species: human

Source: synthetic

Quantity: 50 µg

Concentration: 2 mg/ml

Formulation: DMSO

Molecular Weight: 9.16 kDa

Purity: >98% by InstantBlue™ SDS-PAGE


Stability/Storage: 12 months at -70°C; aliquot as required

Protein Sequence:

MQIFVKLTGKTTITLEVEPSDTIEN

VKAKIQDKEGIPDPQQLIFAGKQL

EDGRTLSDYNIQKESLHLVLRRLRG



Ubiquitin (amino acid residues 1-76)
C-terminally tagged
with 5-TAMRA-Lys(Ub)-Gly-OH
Accession number: P62987

Quality Assurance


Purity:

4-12% gradient SDS-PAGE

InstantBlue™ staining

Lane 1: MW markers

Lane 2: 1 µg Ubiquitin-Lys-TAMRA



Protein Identification:

Confirmed by mass spectrometry.

Activity Assay:

The activity of 5-TAMRA-Lys(Ub)-Gly-OH was validated by determining the decrease in mP (measured at Excitation 540, Emission 590) resulting from cleavage of the fluorophore (TAMRA) from Ubiquitin after incubation with UCHL3 (deubiquitylase). UCHL3 was incubated with 5-TAMRA-Lys(Ub)-Gly-OH and fluorescence intensities were measured in the S (parallel) and P (perpendicular) at four time points (0min, 30min, 60min and 90min), from this data mP values were calculated.



www.ubiquigent.com
Dundee, Scotland, UK

ORDERS / SALES SUPPORT

International: +1-617-245-0003
US Toll-Free: 1-888-4E1E2E3 (1-888-431-3233)
Email: sales.support@ubiquigent.com

UK HQ and TECHNICAL SUPPORT

International: +44 (0) 1382 381147 (9AM-5PM UTC)
US/Canada: +1-617-245-0020 (9AM-5PM UTC)
Email: tech.support@ubiquigent.com

Email services@ubiquigent.com for enquiries regarding compound profiling and/or custom assay development services.

© Ubiquigent 2012. Unless otherwise noted, Ubiquigent, Ubiquigent logo and all other trademarks are the property of Ubiquigent, Ltd.

Limited Terms of Use: For research use only. Not for use in humans or for diagnostics. Not for distribution or resale in any form, modification or derivative OR for use in providing services to a third party (e.g. screening or profiling) without the written permission of Ubiquigent, Ltd.

Lot-specific COA version tracker: v1.0.0