UBE2A (HR6A) [6His-tagged]

E2 – Ubiquitin Conjugating Enzyme

Alternate Names: HHR6A, HR6A, RAD6A, UBC2, EC 6.3.2.19, Ubiquitin-conjugating enzyme E2A

Cat. No. 62-0071-020 Quantity: 20 µg Lot. No. 1822 Storage: -70°C

FOR RESEARCH USE ONLY NOT FOR USE IN HUMANS



CERTIFICATE OF ANALYSIS Page 1 of 2

Background

The enzymes of the ubiquitylation pathway play a pivotal role in a number of cellular processes including the regulated and targeted proteasomal degradation of substrate proteins. Three classes of enzymes are involved in the process of ubiquitylation; activating enzymes (E1s), conjugating enzymes (E2s) and protein ligases (E3s). UBE2A is a member of the E2 conjugating enzyme family and cloning of the human gene was first described by Koken et al. (1991). UBE2A shares 70% identity with its veast homologue but lacks the acidic C-terminal domain. The ring finger proteins RAD5 and RAD18 interact with UBE2A and other members of the RAD6 pathway (Ulrich and Jentsch, 2000). Phosphorylation of UBE2A by CDK1 and 2 increases its activity during the G2/M phase of the cell cycle (Sarcevic et al., 2002). UBE2A is reguired for post-replicative DNA damage repair in eukaryotic cells and it is thought binding to ZNF198 may be involved in this process (Kunapuli et al., 2003). A nonsense mutation resulting in the loss of a 25 amino acid region in the C-terminal domain of UBE2A has been identified as a cause of a novel X-linked mental retardation (XLMR) syndrome (Nascimento et al., 2006).

Continued on page 2

Physical Characteristics

Species: human

Source: E. coli expression

Quantity: 20 µg

Concentration: 1 mg/ml

Formulation: 50 mM HEPES pH 7.5. 150 mM sodium chloride, 2 mM dithiothreitol, 10% glycerol

Molecular Weight: ~21 kDa

Purity: >98% by InstantBlue™ SDS-PAGE

Stability/Storage: 12 months at -70°C;

aliquot as required

Protein Sequence:

MGSSHHHHHHSSGLVPRGSHMASMTG GQQMGRDPNSSSVD**S**TPARRRLMRD FKRLQEDPPAGVSGAPSENNIMVW NAVIFGPEGTPFEDGTFKLTIEFT EEYPNKPPTVRFVSKMFHPNVY ADGSICLDILQNRWSPTYDVSSILT SIQSLLDEPNPNSPANSQAAQLYQENK REYEKRVSAIVEQSWRDC


Tag (bold text): N-terminal His Protease cleavage site: Thrombin (LVPR ▼GS) UBE2A (regular text): Start bold italics (amino acid residues

Accession number: NP_003327

Quality Assurance

Purity:

4-12% gradient SDS-PAGE InstantBlue™ staining Lane 1: MW markers Lane 2: 1 µg His-UBE2A



Protein Identification:

Confirmed by mass spectrometry.

E2-Ubiquitin Thioester Loading Assay:

The activity of His-UBE2A was validated by loading E1 UBE1 activated ubiquitin onto the active cysteine of the His-UBE2A E2 enzyme via a transthiolation reaction. Incubation of the UBE1 and His-UBE2A enzymes in the presence of ubiquitin and ATP at 30°C was compared at two time points, T₀ and T₁₀ minutes. Sensitivity of the ubiquitin/His-UBE2A thioester bond to the reducing agent DTT was confirmed.



Dundee, Scotland, UK

ORDERS / SALES SUPPORT

International: +1-617-245-0003

US Toll-Free: 1-888-4E1E2E3 (1-888-431-3233) Email: sales.support@ubiquigent.com

UK HQ and TECHNICAL SUPPORT

International: +44 (0) 1382 381147 (9AM-5PM UTC) US/Canada: +1-617-245-0020 (9AM-5PM UTC) Email: tech.support@ubiquigent.com

Email services@ubiquigent.com for enquiries regarding compound profiling and/or custom assay development services.

© **Ubiquigent 2011**. Unless otherwise noted, Ubiquigent, Ubiquigent logo and all other trademarks are the property of Ubiquigent, Ltd.

Limited Terms of Use: For research use only. Not for use in humans or for diagnostics. Not for distribution or resale in any form, modification or derivative OR for use in providing services to a third party (e.g. screening or profiling) without the written permission of Ubiquigent, Ltd.

Lot-specific COA version tracker: v1.0.0

UBE2A (HR6A) [6His-tagged]

E2 - Ubiquitin Conjugating Enzyme

Alternate Names: HHR6A, HR6A, RAD6A, UBC2, EC 6.3.2.19,

Ubiquitin-conjugating enzyme E2A

 Cat. No.
 62-0071-020
 Quantity:
 20 μg

 Lot. No.
 1822
 Storage:
 -70 °C

FOR RESEARCH USE ONLY NOT FOR USE IN HUMANS



CERTIFICATE OF ANALYSIS Page 2 of 2

Background

Continued from page 1

References:

Koken MH, Reynolds P, Jaspers-Dekker I, Prakash L, Prakash S, Bootsma D, Hoeijmakers JH (1991) Structural and functional conservation of two human homologs of the yeast DNA repair gene RAD6. *Proc Natl Acad Sci U S A* **88**, 8865-9.

Kunapuli P, Somerville R, Still IH, Cowell JK (2003) ZNF198 protein, involved in rearrangement in myeloproliferative disease, forms complexes with the DNA repair-associated HHR6A/6B and RAD18 proteins. *Oncogene* 22, 3417-23.

Nascimento RM, Otto PA, de Brouwer AP, Vianna-Morgante AM (2006) UBEZA, which encodes a ubiquitin-conjugating enzyme, is mutated in a novel X-linked mental retardation syndrome. Am *J Hum Genet* **79**, 549-55.

Sarcevic B, Mawson A, Baker RT, Sutherland RL (2002) Regulation of the ubiquitin-conjugating enzyme hHR6A by CDK-mediated phosphorylation. *EMBO J* 21, 2009-18.

Ulrich HD, Jentsch S (2000) Two RING finger proteins mediate cooperation between ubiquitin-conjugating enzymes in DNA repair. *EMBO J* **19**, 3388-97.



Dundee, Scotland, UK

ORDERS / SALES SUPPORT

International: +1-617-245-0003

US Toll-Free: 1-888-4E1E2E3 (1-888-431-3233) Email: sales.support@ubiquigent.com

UK HQ and TECHNICAL SUPPORT

International: +44 (0) 1382 381147 (9AM-5PM UTC) US/Canada: +1-617-245-0020 (9AM-5PM UTC)

Email: tech.support@ubiquigent.com

Email services@ubiquigent.com for enquiries regarding compound profiling and/or custom assay development services.

© **Ubiquigent 2011**. Unless otherwise noted, Ubiquigent, Ubiquigent logo and all other trademarks are the property of Ubiquigent, Ltd.

Limited Terms of Use: For research use only. Not for use in humans or for diagnostics. Not for distribution or resale in any form, modification or derivative OR for use in providing services to a third party (e.g. screening or profiling) without the written permission of Ubiquigent, Ltd.

Lot-specific COA version tracker: v1.0.0